


LIGO Highlights On behalf of Korean LIGO members

Kyung-ha Lee

Sungkyunkwan University khlee54@skku.edu

2025 Particle/Nuclear/Astro Workshop

November 19, 2025





GW Talks

- 1. LIGO Highlight (NOW): Kyung-ha Lee
- 2. GW Universe (5pm Today): Hyung Mok Lee
- 3. KGWG Highlight (5pm Friday): Hyung Won Lee
- 4. YEMIGO (12pm Saturday): John Oh



Overview

1. LIGO Introduction


- Observing Plans
- Structure of LSC
- Korean LIGO Group Summary

2. LIGO Experimental Work in Korea

- Coating Brownian Noise
- 3. LIGO Data Analysis Work in Korea
 - CBC & DetChar & etc
- 4. Summary & Future Work




LIGO Observing Plan






LIGO Observing Plan





LIGO Observing Plan



- Observing Plans are constantly changing.
- O4 was just concluded last night!
- Upgrading & Commissioning period started now!
- Partners: IceCube, CHIME, IPN, Swift, Fermi, etc



LIGO Scientific Collaboration (LSC)

- Experimental
 - Optics Working Group
 - KGWG
 - Advanced Interferometer
 Configurations Group
 - Newtonian Noise Group
 - Seismic Isolation & Suspension Working Group
 - Controls Group
 - Lasers and Auxiliary Optics
 Group
 - Quantum Noise Group

- ◆ Data Analysis & Theory
 - Burst Search Group
 - KGWG
 - Compact Binary
 Coalescence Search Group
 - KGWG
 - Continuous Wave Search Group
 - Stochastic Search Group
 - Detector Characterization
 Working Group
 - KGWG
 - Data Analysis Council



LIGO/KGWG

- Members 16 [2024 Jan.] (-> 19 [current, three students were added])
 - 8: Permanent Members (1 SNU, 1 Ewha, 2 NIMS, 1 Inje, 1 KISTI, 1 Chungang, 1 SKKU)
 - 4 Researchers (1 SNU, 1 Ewha, 1 UNIST, 1 KASI)
 - 7 Students (1 NIMS, 1 Ewha, 1 ChungAng, 4 SKKU)
 - Authors: 13



Recent Activities of KGWG in LSC (2024-2025)

Software

- Investigation of the effects of limited number of bins on the parameter estimation, for further development of parameter estimation code,
- Development of fast templates for microlensed events using surrogate modelling
- Review of the results from tidal parameter estimation and EoS inference pipelines for LVK publications based upon nuclear physics perspectives, i.e. the effect of nuclear symmetry energy.
- Optimization of parameter estimation for Compact Binary Coalescences with loweccentricity

Waveforms

- Support for precession in gw_eccentricity, review the code and add to PE post processing.
- Provide NR waveforms for Black Hole Capture Events



Recent Activities of KGWG in LSC (2024-2025)

Data Analysis

- Search for BH capture events with Coherent Wave Burst (cWB) + Gaussian Mixture Model (GMM) pipelines
- Simulation-based study for finding multi-messenger signals of strong lensing utilizing EM catalogs
- Offline PE activities including PE rota for O4
- Analysis of low-mass CBCs from O4 and/or O3 using TaylorF2Ecc, as contribution for O4 fullauthor paper

Review

- Chairing analysis review for the S230814ah paper (GW230814, the loudest event so far)
- Detector Characterization
 - Maintain and update the codes of machine learning model (MSNN, explainable AI) to classify and trace the responsible or witnessing channels of known noiseCoating
 - Monitoring known or new Data Quality issues in the LIGO interferometers



Recent Activities of KGWG in LSC (2024-2025)

- Shifts
 - DQ Shifts
 - Rapid Response Team (RRT) Shift
- Computing Resources
 - Managing Condor on KISTI Cluster for IGWN
 - Maintaining KISTI Cluster (Tier 2 Center of LIGO Data Grid) H/W for IGWN
- Public Outreach
 - Maintain and refine of the portable GW detection simulator for exhibitions at educational and outreach events.
- Experiments
 - Coating candidate material search



Potential Collaborations with us!


- ◆ Data analysis
 - Multi-messenger astrophysics
 - Simulations between LIGO & nuclear physics / high energy physics
 - Theory?
- Experimental
 - Each noise needs various research expertise.
 - Coatings —> Optics & Material science
 - Quantum —> Optics & Quantum
 - Hardware contribution is essential for LIGO.
- ◆ Becoming LIGO member might not be very straightforward, but LSC is definitely welcoming all multi-messenger collaborators & experimental expertise who can contribute to upgrade plans!



LIGO Detector Introduction





LIGO Detector Introduction: Interferometer 101





LIGO Detector Introduction: Noise Curve






 \mathbb{C}_{M} \exists \exists M \exists M



LIGO Detector Introduction: Noise Curve






LIGO Experimental Work in Korea




LIGO Experimental Work in Korea Suspension Thermal & Coating Thermal







Suspension Thermal Noise: Quadruple Monolithic Suspension





Coating Brownian Noise: Why Important?






Plots from LIGO-T15TBI-v1, I.S. White Paper LIGO-T1800042-v5 A+ Design Curve


Coating Brownian Noise: Current aLIGO Coating

- Ion-beam sputtered (IBS) dielectric multi-layer amorphous coatings
- Materials: SiO₂ (low index) + Ti:Ta₂O₅ (high index)



laser beam radius coating mechanical loss

- Optical specs:
 - Diameter: 34 cm
 - Optical Absorption: ~0.5 ppm
 - Micro-roughness scatter: ~5 ppm
 - Uniformity: ~0.5 nm





Material	Refractive index	Loss
Silica SiO ₂	1.45	0.4 x 10 ⁻⁴
Tantala Ta ₂ O ₅	2.03	3.4 x 10 ⁻⁴
Titania-doped tantala Ta ₂ O ₅ - TiO ₂	2.07	2.3 x 10 ⁻⁴




Coating Brownian Noise: Amorphous + Dopant + Annealing

- Why amorphous?
 - Loss peak at low temperature low RT loss
 - Less scattering
- Why dopants?
 - Can reduce RT mechanical loss
 - Can increase crystallization temperature
- Why annealing?
 - Can reduce RT mechanical loss











Coating Brownian Noise: A+LIGO Coatings Research Model

How to find a new coating material?






Coating Brownian Noise: A+LIGO Coatings Research Model






Coating Brownian Noise: A+LIGO Coatings Research Model






Coating Brownian Noise: Atomic Structure Characterization


- Characterization of structure of amorphous materials more difficult than crystalline
 - PDF (Pair Distribution Function) Analysis:
 - Most efficient to get SRO & Intermediate SRO information
 - GIPDF (Grazing Incidence PDF) from SLAC SSRL (Stanford Linear Accelerator Center, Stanford Synchrotron Radiation Lightsource)
 - ePDF (electron PDF) via TEM (Transmission Electron Microscope)
 - FEM (Fluctuation Electron Microscopy) Analysis: €
 - Conducted @ Lawrence Berkeley National Lab, USA
 - Most efficient to get MRO information









Coating Brownian Noise: What kind of atomic structure?







Corner-sharing


Edge-sharing

Face-sharing

Three types of polyhedral connections exist.




Coating Brownian Noise: What kind of atomic structure?



Corner-, edge- and face- sharing polyhedra register distinct peaks in measured PDFs.



Coating Brownian Noise: PDF Results: Material Comparison



Corner-shared polyhedra:


Silica* > Tantala > Zirconia-doped tantala > Silica-doped hafnia (*silica is not in the plot but is known to be corner-shared)


Room temperature mechanical loss:

Trends in corner-shared polyhedra and loss are flipped*.
*RT loss for silica-doped hafnia is not measured.




Coating Brownian Noise: **Energy Dispersive Spectroscopy (EDS)**






Position [um]



- Optimal coating for A+LIGO upgrade: 52 layer of dielectric HR coating
 - 44% TiO₂-doped GeO₂ / SiO₂
 - 600°C, 100hr annealing
- Discrepancy between expected CTN from single-layer vs actual multi-layer coating.
 - → Expected: 0.5CTN @ aLIGO
- Possible source: interfaces between TiGe/SiO₂ layers Minhyo Kim, 2024 LVK Meeting




Coating Brownian Noise: Energy Dispersive Spectroscopy (EDS)



https://www.gatan.com/techniques/eds

- EDS: Captures x-rays from excited atoms inside of material
- EDS can show atomic composition mapping at the interface.
 - → Relative, qualitative comparison





LIGO Data Analysis Work in Korea



Korean LIGO Group Summary


- ◆ 13 authors (up to 2024-2025 MOU), 16 members total.
 - Hyungmok Lee (SNU): Management, election administration, RRT shifts
 - Gungwon Kang (CAU): Burst pipeline, outreach, RRT shifts
 - Jiyoon Sun (CAU): Burst pipeline, offline search of burst signals
 - Young-min Kim (KASI): DetChar Code development, Nuclear & High Energy collaboration
 - Kyungmin Kim (KASI): O4 lensing paper, multi-messenger signals of lensing
 - Jeongcho Kim (KASI): Parameter estimations, evaluation and development of codes
 - Chunglee Kim (EWU): Parameter estimations, evaluation of methods, RRT shifts
 - Kyung-ha Lee (SKKU): Structural studies of coatings, material evaluation & assessments
 - Minhyo Kim (SKKU): Structural studies of coatings, material evaluation & assessments
 - Jeonghyeon Ju (SKKU): Structural studies of coatings, material evaluation & assessments
 - Sunjae Lee (SKKU): Structural studies of coatings, material evaluation & assessments
 - Sangwook Bae (KISTI): Maintaining & managing KISTI cluster
 - Md Shaikh (SNU): Eccentric waveform models for CBC systems
 - Whansun Kim (NIMS) (Not LIGO author): Developing new ML tool for DetChar
 - Byeol Kim (NIMS) (Not LIGO author): Monitoring DQ issues
 - Hyungwon Lee (IJU) (Not LIGO author): Parameter estimations, evaluation&development of codes



Korean LIGO Group Summary

Strong lensing of dark sirens and galaxies as a probe of the Hubble constant [in preparation]


- Motivation
 - Simultaneous observations of GW170817 & GRB170817 → GWs can be a probe of the Hubble constant, H0!
 - o GW events without identified EM counterparts, a.k.a. dark sirens, can also be such a probe.
 - Accurate sky localization and precise estimation of luminosity distance to the source of dark sirens can be achieved if the events are strongly lensed.
- Methods
 - Simulated dark sirens and simulated galaxy catalog
 - Single lensed dark sirens w/ the Hubble-Lemaître law
 - Multiple lensed dark sirens w/ combining individual inferences of H0
- Result (preliminary)
 - Using quadruple lensed images is better than double images in
 - accurately localizing the host galaxy of a dark siren.
 - well constraining and precisely inferring H0.
- Contact: Kyungmin Kim (KASI; kkim@kasi.re.kr)







Parameter Estimation Group (1/2)

- Simulation of parameter estimation for binary inspiral with aligned spin and eccentricity
- Waveforms: TaylorF2Ecc-Inspiral
- Advanced LIGO and Virgo sensitivity








Assuming GW20015 parameters for mass and distance (NS-BH) with e0=0.2, eccentricity is more precisely constrained for higher aligned spin magnitude



Parameter Estimation Group (2/2)

- Comparison of two pipelines: lalinference_mcmc vs Bilby
- Purpose: to investigate the accuracy improvements for important parameters such as distance and mass of the sources with external constraints on viewing angle through EM observations (BNS of NSBH)
- There are some differences in estimated parameters for different pipelines: We need more thorough tests



Gaussian Mixture Modelling(GMM)

슬라이드 제공: 선지윤


- Describes a non-Gaussian distribution of data as a superposition of multivariate $\operatorname{Gaus}_{p(\vec{x_i})} = \sum_{j=1}^K w_j \mathcal{N}(\vec{x_i} | \mu_j, \Sigma_j)$
- Model parameters:

$$\hat{\boldsymbol{\mu}}_{k} = \frac{\sum_{i=1}^{n} r_{ik} \mathbf{x}_{i}}{N_{k}}$$

$$\hat{\Sigma}_{k} = \frac{1}{N_{k}} \sum_{i=1}^{n} r_{ik} \left(\mathbf{x}_{i} - \boldsymbol{\mu}_{k}\right) \left(\mathbf{x}_{i} - \boldsymbol{\mu}_{k}\right)^{T}$$

$$\hat{w}_{k} = \frac{\sum_{i=1}^{n} r_{ik}}{n} = \frac{N_{k}}{n}.$$

For kth
Gaussian
component





$$\ln \mathcal{L} = \sum_{i=1}^{n} \ln p(\vec{x}_i|\Theta) = \sum_{i=1}^{n} \ln \left\{ \sum_{j=1}^{K} w_j N(\vec{x}_i|\mu_j, \Sigma_j) \right\}$$


GW detections: Application to O3 data

슬라이드 제공: 선지윤

List of detected events:

Event Name	Т	GMM+ iFAR (yr)
GW190412	18.46	85.93
$ m GW190519_153544^{-2}$	7.94	19.09
$GW190828_063405$	5.79	15.62
$GW200224_222234$	10.74	3.39
$GW190706_222641$	-0.14	1.75
$GW190521_074359$	-0.5	1.43

 No non-CBC events were found during the O3 run



PhysRevD.110.083032 (L. Smith, S. Ghosh, J. Sun et al.)

- GMM brings sensitivity improvements for many of the tested waveforms.
 - Especially, GMM has noticeable sensitivity for GAs and CSs, and it proves that GMM can
 - effectively mitigate blip glitches
- First case of cWB flavours to have better sensitivity on H1L1V1 network than on H1L1 network
- Further investigations ...
 - Search for optimal setting of GMM configuration
 - Test on released regulators

Slide from Hyung Mok Lee's presentation (LSC Report 2025)



LIGO/KAGRA DECHAR Group

- 1. CAGMon and EEMD **김별** (NIMS), 오정근(NIMS), 김영민 (KASI)
- 2. MSNN and Occlusion sensitivity 김환선(NIMS), 김영민(KASI)
- 3. t-SNE & PCA analysis (unsupervised learning) for classification **이우진**, 곽규진, 정기현 (UNIST)
- 4. KAGRA O4a hveto analysis **정기현**, Nursyifa Ruhama, 곽규진 (UNIST)
- 5. Bayesian Analysis for NS EoS 김영민 (KASI)



Summary & Future Work

- ✓ LIGO has successfully running to detect gravitational waves but noise sources has been the main constraint for the sensitivity.
- ✓ Instrumental R&D is the core research to overcome that challenge.
- ✓ Collaborations established working groups assigned for each noise source for high-level expertise.
- √ Korean groups are expanding their research field from data analysis to experimental R&D.



Summary & Future Work

- ✓ LIGO has successfully running to detect gravitational waves but noise sources has been the main constraint for the sensitivity.
- ✓ Instrumental R&D is the core research to overcome that challenge.
- ✓ Collaborations established working groups assigned for each noise source for high-level expertise.
- √ Korean groups are expanding their research field from data analysis to experimental R&D.

Gravitational wave field has JUST started its first baby step, and more exciting things are to come! :)



Thank you for your attention.

Kyung-ha Lee khlee54@skku.edu







