



IceCube Neutrino Observatory





THE ICECUBE COLLABORATION

CANADA

Universiteit Gent

UCLouvain

Queen's University University of Alberta–Edmonton

Université libre de Bruxelles

Vrije Universiteit Brussel

DENMARK

University of Copenhagen

GERMANY

Deutsches Elektronen-Synchrotron ECAP, Universität Erlangen-Nürnberg Humboldt–Universität zu Berlin Karlsruhe Institute of Technology Ruhr-Universität Bochum **RWTH Aachen University** Technische Universität Dortmund Technische Universität München Universität Mainz Universität Wuppertal Westfälische Wilhelms-Universität Münster

ITALY

University of Padova

JAPAN

Chiba University

NEW ZEALAND

University of Canterbury

REPUBLIC OF KOREA

Chung-Ang University Sungkyunkwan University

SWEDEN

Stockholms universitet Uppsala universitet

SWITZERLAND Université de Genève

* TAIWAN

Academia Sinica

UNITED KINGDOM University of Oxford

UNITED STATES

Clark Atlanta University Columbia University **Drexel University** Georgia Institute of Technology Harvard University Lawrence Berkeley National Lab Loyola University Chicago Marquette University

Massachusetts Institute of Technology Mercer University Michigan State University Ohio State University Pennsylvania State University South Dakota School of Mines and Technology Southern University and A&M College **Stony Brook University** University of Alabama University of Alaska Anchorage University of California, Berkeley University of California, Irvine University of Delaware University of Kansas

University of Maryland University of Nevada, Las Vegas University of Rochester University of Texas at Arlington University of Utah University of Wisconsin–Madison University of Wisconsin-River Falls **Yale University**

FUNDING AGENCIES


Fonds de la Recherche Scientifique (FRS-FNRS) Fonds Wetenschappelijk Onderzoek-Vlaanderen (FWO-Vlaanderen)


German Research Foundation (DFG) Deutsches Elektronen-Synchrotron (DESY)

Federal Ministry of Education and Research (BMBF) Japan Society for the Promotion of Science (JSPS) Knut and Alice Wallenberg Foundation Swedish Polar Research Secretariat

The Swedish Research Council (VR) University of Wisconsin Alumni Research Foundation (WARF) US National Science Foundation (NSF)







Cherenkov Detector using 1 km³ South Pole ice


IceCube Science

- Neutrino Astrophysics
 - Diffuse Neutrinos
 - Search for Neutrino source
 - Multi-messenger astrophysics
- **Neutrino particle physics**
- Neutrino oscillations
- Sterile neutrinos & NSI search
- Cosmic Ray physics
 - Cosmic ray anisotropy
 - Components of Cosmic ray
 - Cosmic ray spectrum
- Dark matter & BSM physics
 - Dark matter indirect detection
- Axion, Magnetic monopole, SUSY.

Event Topology (Track vs. Cascade)


Track-like

- Good angular resolution: 0.5° for E > 10 TeV
 → prime channel for point sources
- Energy resolution factor ~ 2




Cascade-like


- Good energy resolution < 10% for $E > 10 \,\mathrm{TeV}$
- Angular resolution $\lesssim 5^{\circ}$ for $E > 50 \,\mathrm{TeV}$ (contained events)




Event Topology (Track vs. Cascade)



$$\nu_e$$
 + N \rightarrow e + X EM shower + Hadron shower (Cascades)






Tau decay (double pulse/shower)

Event Selections


Northern sky through-going Track analysis

- -Effective Volume is large (more events)
- -Neutrino Search only northern sky (2π)
- -Energy measurement in a sampling way




Veto-based analysis

- -Veto-Fiducial volume division
- -Effective Volume is small (less events)
- -All sky Neutrino Search (4π)
- -Events can contain cascades.
- -Calorimetric energy measurement




Neutrino Flux (Astrophysical diffuse neutrino)




Astrophysical Neutrino Flux


Northern sky through-going Track analysis



All sky Cascades analysis



Astrophysical Neutrino Flux



Neutrino Energy [GeV]


$-2\Delta \ln \mathscr{L}$ compared to SPL
0
1.8 (0.9σ)
19.8 (4.2σ)
22.3 (3.9σ)
27.3 (4.7σ)

Astrophysical Neutrino Flux is not a single mechanism!

Neutrino Energy [GeV]

Single Power model is rejected at 4.7σ from Broken Power model (hard at low-E and soft at high-E)


Neutrino Flux (GZK neutrinos, Extremely High Energy)




THE RECORD NEUTRINOS 700 m NEUTRINO With no electric charge, being extremely light and travelling almost at the speed of light, these elementary particles interact only weakly, and therefore very rarely, with matter. Their elusiveness makes them valuable cosmic messengers, able to bring us unique information about the distant universe. On 13 February 2023, at a depth of 3450 metres off the coast of Sicily, in Italy, the ARCA detector of the KM3NeT submarine neutrino telescope recorded an extraordinary THE EVENT DISPLAY signal: produced by a neutrino with a record energy of about 220 PeV, corresponding to 220 million billion electronvolts. This signal, named KM3-230213A, provides the first A view of the KM3-230213A signal evidence that neutrinos with such extreme energies exist in the universe. detected by KM3NeT. The spheres are coloured according to 1600 the detection time and the reconstructed track of the particle is shown. The size of 1400 the blue cone gives an indication on the KM3-230213A IDENTIKIT 1200 amplitude of the signal. 1000 The cosmic neutrino plunged into the Mediterranean Sea and crossed the Malta 800 continental shelf with an inclination of 0.6° above the horizon. During this journey, it travelled almost at the speed of light and interacted with an atomic nucleus, 600 generating an ultra-relativistic muon, which crossed the whole detector. 200 TIME (ns) SICILY THE SOURCES The origin of the ultra-high energy neutrino could have been one of the cataclysmic events that animate our universe, such as an active galactic nucleus or a gamma-ray burst. Or it could be a neutrino generated by the interaction of an ultra-high energy cosmic-ray particle with the cosmic background radiation that KM3NeT Coll, Nature 638 (2025) 8050 permeates the universe. **KM3NeT** 12

IceCube EHE neutrino search & KM3-230213A


$$p_{UHECR} + \gamma_{CMB} \rightarrow \Delta^+ \rightarrow n + \pi^+$$





- -KM3NeT saw EHE candidate at 220 PeV (E^{-2})
- —The same flux would require lceCube to have 70 events but none seen. (> 10σ)
- -Flux uncertainty can give reconciliation among KM3NeT,

Auger, IceCube at most 3σ tension




KM3NeT Coll, Nature 638 (2025) 8050


(Personal) Source Questions

Once in a lifetime (1/70 years)

- IceCube (complete, $1 \text{ } km^3$) 14 years VS ARCA (partial, $0.15 \text{ } km^3$) 1 year
- IceCube + AMANDA + Baikal + Antares + Auger + TA + RADIOs VS ARCA
- If this is a GZK neutrino, why only the highest energy? Lower energies?
- Is it possible to have an astrophysical object with ultra-high-energy neutrino emission only without gamma or EM counter parts?
- If it is a once-in-a-lifetime event, why do we build a new detector?




NGC 1068




- Barred Spiral Galaxy
- 47 million light year away
- Supermassive Black Hole hidden by a cloud of dust.

Cosmic-ray interactions occur in a cloud. Neutrinos are generated in the surrounding medium where photons attenuate.



- New analysis with 13 years of data.
- The hottest spot is closer to NGC 1068.
- Significance : 5.0σ local, 4.0σ global
- Prefers softer spectrum ($\sim E^{-3.4}$), indication of a more complex power law.


Observation of high-energy neutrinos from the Galactic plane





Science, 380, 6652 (2023)

Ice, Ice, South Pole Ice is very complicated Cherenkov medium


Most of IceCube analysis systematics comes from optical properties of Ice







- South Pole Ice is very clear. However, it has a layered structure which shows different optical characteristics (scattering and absorption).
- Ice flows and light travel in birefringent way (along the flow direction; SE->NW).
- Refrozen (Hole) Ice is different from the bulk ice.




The IceCube Upgrade


- Instrument additional 7 strings
 - 60-100 DOMs in each string
- Detect neutrinos below 10 GeV
- Construction began and full deployment in 2025/2026
- Testbed for IceCube-Gen2

- Neutrino Oscillations study,
- Neutrino Mass Hierarchy measurement with the Upgrade.
- Neutrinos with Earth matter and density effect in varying baseline
 - Understanding the systematics is the key to the measurement
 - Additional calibration devices is going to be added











The IceCube Upgrade Contributions




Upgrade Scope (10 km view)

- Major Parts of Detector Instrumentation Instrumentation (*largely in-kind* contributions)
 - 277 D-Eggs (+ spares) (Chiba University)
 - 402 mDoms (+ spares) (Germany; 50% built at DESY-Zeuthen, 50% built at Michigan State University)
 - 7 Main Cable assemblies (Michigan State University) / Surface cables (Sweden)
 - In optical module camera systems (SKKU)
 - Calibraton Devices / Special Devices (Various inkind contributors)
- Drill and deploy 7 strings of detector elements
 - Drill (University of Wisconsin, Physical Sciences Laboratory)
 - Logistics (NSF Office of Polar Programs)


The IceCube Upgrade Contributions





























Atmospheric Neutrino Oscillation Physics Sensitivity

The latest atmospheric neutrino oscillation result (DeepCore)




The DeepCore+Upgrade Sensitivity



The Upgrade will enhance oscillations sensitivity significantly (i.e. $\sin^2\theta_{23}\sim4\%$)

Atmospheric Neutrino Oscillation Physics Sensitivity

True Normal Ordering



The Upgrade will enhance oscillations sensitivity significantly with increased statistics and improved systematics

Korea IceCube Activities

IceCube @ Chung-Ang University


- Study atmospheric neutrinos. Especially electron neutrinos
- Participation in the IceCube-Upgrade simulations. i.e. Photon Propagation & mDOM response.
- Develop a low-energy reconstruction tools using Transformer-based neural network



IceCube @ Sungkyunkwan University


- Muon-track reconstruction using a neural network that fits Gupta– Kundu–based arrival-time PDFs
- Generative model for IceCube neutrino event simulation and end-toend reconstruction using flow and diffusion model
- Optimize pulse cleaning and upgrade PMT noise model based on its timing and charge characteristics





Track Reconstruction (Gupta-Kundu-based time PDF) at SKKU

• For a given track $\vec{\theta}$, the geometric arrival time t_{res} of a Cherenkov photon is as follows:



$$t_{\text{geo}} = t_0 + \frac{\hat{\mathbf{p}} \cdot (\mathbf{r_i} - \mathbf{r_0}) + d \tan \theta_c}{c_{\text{vac}}}$$

• Due to the optical properties of the detector medium, the arrival time of a Cherenkov photon is stochastically delayed. The delay, t_{res} , is modeled by the **Pandel function**.


$$p(t_{\text{res}}, | d, \vec{\theta}) = \frac{\beta^{\alpha}}{\Gamma(\alpha)} t_{\text{res}}^{\alpha-1} \exp(-\beta t_{\text{tes}})$$

• The track is obtained by maximizing the following - form known as the MPE likelihood


$$\mathscr{L}_{\text{MPE}}(\vec{\theta}) = \prod_{i}^{\text{1st hits}} n_i \cdot p(t_{\text{res},i} | d, \vec{\theta}) \cdot \left(\int_{t_{\text{res},i}}^{\infty} p(t | d, \vec{\theta}) dt \right)^{n_i - 1}$$

Replace the Pandel Function with Gupta-Kundu Generalized Exponential function using neural network training.

Pulse Cleaning: Modularization and **Optimization (SKKU)**









Before Cleaning

New cleaning segment




IceCube Upgrade event reconstruction

Transformer-based neutrino event reconstructions (Aachen-CAU)



IceCube Upgrade Simulation development at CAU



Separation of a code for GPU photon propagation in ice from Geant4 neutrino production makes Upgrade simulation chain more flexible and accessible.


Atmospheric Neutrino Oscillation Physics at CAU

New Veto Algorithms (VICH) by CAU Towards the Upgrade ν_e appearance analysis

 $4 \text{ GeV} < \text{Energy} < 15.8 \text{ GeV} \& 120^{\circ} < \theta_{\text{zen}} < 180^{\circ}$ 10^{-1} ¥ 1.5


It is clear that reconstruction and PID will be two challenges in the Upgrade. CAU will work on those in $\nu_{\mu} \to \nu_{e}$ analysis

particle speed




IceCube-Gen2






IceCube-Gen2: Eyes to the Violent Cosmos

J. Phys. G: Nucl. Part. Phys. 48 060501 (2021)



Neutrinos from TeV to EeV
Cosmic particle acceleration
Sources and propagation of HE particles

Fundamental physics with HE neutrinos



Summary and Outlook


The IceCube Upgrade is starting now


- Korean IceCube members (~8) are growing. CAU & SKKU
- We are interested in Physics analyses in the Upgrade.
- Participate in detector response simulations and event reconstructions.
- The Upgrade will happen starting now.
- Korean IceCubers will participate in the Upgrade activities.
- •아이스큐브는 활동성 은하핵 과 같은 천체에서 나오는 중성미자 입자를 탐색하는 실험입니다. 국내에서 이제 출발하지만 SKKU-노창동, CAU-하창현 그룹이 새로운미래를 열것으로 기대하고 있습니다. 많은 관심과 지원 부탁드립니다.



Back-up

Multi-messenger observations of a flaring blazar coincident with high-energy neutrino IceCube-170922A







Science 361, eaat1378 (2018)

IceCube neutrino alert prompted γ -ray follow-up searches by e.g. FermiLAT & MAGIC, resulting in flaring blazar TXS 0506+056, 0.1 deg. from the neutrino.


Neutrino emission from the direction of the blazar TXS 0506+056 in association with the IceCube-170922A



Galactic Plane: Charged Pions must exist; Neutrinos should be there



Excess over the Galactic Plane : 4.71 σ , 4.37 σ , and 3.96 σ depending on models, rejecting the background-only hypothesis.




Trial factor correction : penalty on p-values —>4.48 σ

Right Ascension α 1.5 2.0 2.5 3.0 3.5 4.0 seemingly hot spots to be fluctuations

Implications:


Science, 380, 6652 (2023)



- KRA models and best fits are consistent, except very high energies.
- π^0 model differs by a factor 5, suspicious of CR propagation or spectral index from GC. Best fit agrees with Tibet gamma ray measurements.
- Neutrinos from the Galactic Plane contribute 6~13% to the astrophysical flux at 30 TeV.

Improved data analysis


Reprocessing of the previous data uniformly with new event reconstruction algorithms and new calibrations.



The application of the latest directional reconstruction

Better energy reconstruction by extracting more accurate DOM charge and using Machine Learning.

Evidence for neutrino emission from the nearby active galaxy NGC 1068



E_v [GeV]